

UNCLASSIFIED

AD 633 718

THE GENERALIZED CAYLEY-HAMILTON THEOREM
IN N DIMENSIONS

John S. Lew

Brown University
Providence, Rhode Island

March 4, 1965

Processed for . . .

DEFENSE DOCUMENTATION CENTER
DEFENSE SUPPLY AGENCY



U. S. DEPARTMENT OF COMMERCE / NATIONAL BUREAU OF STANDARDS / INSTITUTE FOR APPLIED TECHNOLOGY

UNCLASSIFIED

NASA Scientific and Technical Information Facility

operated for the National Aeronautics and Space Administration by Documentation Incorporated

Post Office Box 33
College Park, Md 20740

Telephone | Area Code 301
| 779-2121

FACILITY CONTROL NO. 23614

DATE 6/25/66

ATTACHED IS A DOCUMENT ON LOAN

NASA Scientific and Technical Information Facility

TO: Defense Documentation Center
Attn: DDC-IRC (Control Branch)
Cameron Station
Alexandria, Va. 22314


In accordance with the NASA-DOD Cooperative AD Number Assignment Agreement it is requested that an AD number be assigned to the attached report.

☒ As this is our only available copy the return of the document (with AD number and any applicable distribution limitations) to the address below is essential.

☐ This document may be retained by DDC. If retained, please indicate AD number and any applicable distribution limitations on the reproduced copy of the title page and return to the address below.

Return Address: NASA Scientific and Technical Information Facility
Attention: INPUT BRANCH
P. O. Box 33
College Park, Maryland 20740

AD 633718



Division
of

APPLIED MATHEMATICS

ONR Technical Report No. 6
ARPA Technical Report No. AM-20

(THE GENERALIZED CAYLEY-HAMILTON THEOREM
IN n DIMENSIONS

by

John S. Lew

BROWN UNIVERSITY

FC# 33614

Providence
Rhode Island

MP

The Generalized Cayley-Hamilton Theorem in n Dimensions

John S. Lew
Brown University

In 1945, Reiner⁽¹⁾, by means of the Cayley-Hamilton theorem, obtained a canonical form for a polynomial relation between a stress matrix and a strain-velocity matrix; since that time an extensive theory has been developed for canonical forms of non-linear constitutive equations. More recently, for a polynomial relation between one tensor and a number of other tensors, the problem of finding the restrictions imposed by a symmetry group was reduced by Smith and Rivlin⁽²⁾, and by Pipkin and Rivlin⁽³⁾, to that of finding an integrity basis for a set of such tensors; and then, for the full (or proper) orthogonal group in Euclidean 2-space or 3-space, such a basis was determined by Rivlin⁽⁴⁾, Spencer and Rivlin⁽⁵⁾, and Spencer⁽⁶⁾, and its irreducibility proven by Smith⁽⁷⁾.

In this development, an important tool has been a generalization of the Cayley-Hamilton theorem, in 2-space or 3-space, from one to several matrix variables⁽⁸⁾. During this time, it has been clear that the corresponding identity in n -space, for any particular n , could be obtained in a finite but discouraging number of steps; however the form of this relation for an arbitrary n has not been given. We shall obtain this form, which is the intuitive generalization of the results in two and three

-
- (1) See Reference 4.
 - (2) See Reference 7.
 - (3) See Reference 3.
 - (4) See Reference 5.
 - (5) See References 9,10,11.
 - (6) See Reference 8.
 - (7) See Reference 6.
 - (8) See Reference 5.

dimensions, and note that, properly expressed, it is a polynomial relation in the given matrices and their traces with all coefficients ± 1 .

For an arbitrary real or complex $n \times n$ matrix A the Cayley-Hamilton theorem states that

$$1) \quad \sum_{i=0}^n (-1)^i A^{n-i} s_i(A) = 0$$

where $A^0 = I$, and $s_i(A)$ is the i 'th symmetric polynomial in the characteristic roots of A . If we let $t_j(A) = \text{tr } A^j$ for $j = 1, 2, \dots$ then the well-known relations⁽⁹⁾

$$s_1 = t_1$$

$$2) \quad 2s_2 = s_1 t_1 - t_2$$

$$3s_3 = s_2 t_1 - s_1 t_2 + t_3$$

and so forth can be solved recursively for each s_i in terms of t_1, \dots, t_i to yield

$$1!s_1 = t_1$$

$$3) \quad 2!s_2 = t_1^2 - t_2$$

$$3!s_3 = t_1^3 - 3t_1 t_2 + 2t_3$$

and so forth. Thus the Cayley-Hamilton theorem can be expressed as a relation in A and the $t_j(A)$.

In one dimension this process gives

$$4) \quad A - I \text{ tr } A = 0$$

which is trivial; and in two dimensions it gives

$$5) \quad A^2 - A \text{ tr } A + \frac{1}{2} I[(\text{tr } A)^2 - \text{tr } A^2] = 0.$$

(9) See p. 9 of Reference 12.

If we apply to this equation the polarization operator d_{BA} , that is, if we replace A by $A + xB$, for a real variable x , and evaluate the derivative in x at the point $x = 0$, then we obtain

$$6) \quad AB + BA - A \operatorname{tr} B - B \operatorname{tr} A + I[\operatorname{tr} A \operatorname{tr} B - \operatorname{tr} AB] = 0$$

which is the generalized identity in two dimensions⁽¹⁰⁾. Note in this result that all permutations of A and B appear, since A and B need not commute, but that the fraction $\frac{1}{2}$ disappears, since scalars commute and traces have cyclical symmetry.

In three dimensions equations (1) and (3) give

$$7) \quad A^3 - A^2 \operatorname{tr} A + \frac{1}{2} A[(\operatorname{tr} A)^2 - \operatorname{tr} A^2] \\ - \frac{1}{6} I[(\operatorname{tr} A)^3 - 3 \operatorname{tr} A \operatorname{tr} A^2 + 2 \operatorname{tr} A^3] = 0$$

applied to which the polarization operator d_{BA} again yields a relation in two variables. However, we desire a completely polarized relation, in which no matrix variable has degree more than unity. Thus if we also apply d_{CA} for another 3×3 matrix C , and let Σ denote the sum over all permutations of (A, B, C) , then we obtain

$$8) \quad 0 = \Sigma ABC - \Sigma AB \operatorname{tr} C + \Sigma A[\operatorname{tr} B \operatorname{tr} C - \operatorname{tr} BC] \\ - I[\operatorname{tr} A \operatorname{tr} B \operatorname{tr} C - \operatorname{tr} A \operatorname{tr} BC - \operatorname{tr} B \operatorname{tr} CA - \operatorname{tr} C \operatorname{tr} AB \\ + \operatorname{tr} ABC + \operatorname{tr} CBA]$$

which is the generalized identity in three dimensions⁽¹¹⁾. Note here again that all fractions disappear by the properties of scalars and traces.

(10) See Reference 5.

(11) See Reference 5.

Now in the derived expression for each s_1 , the terms correspond to partitions of i , that is, to sequences $\mu = (m_1, m_2, \dots)$ of non-negative integers with $q(\mu) = i$, where

$$9) \quad p(\mu) = \sum_{j=1}^{\infty} (j-1)m_j, \quad q(\mu) = \sum_{j=1}^{\infty} jm_j.$$

Each m_j is interpreted as the number of subsets containing precisely j elements in the corresponding subdivision of a set containing precisely i elements, so that clearly $m_j = 0$ for $j > i$ and thus such sequences have all entries but a finite number equal to zero. If we let τ denote the sequence (t_1, t_2, \dots) with $t_j = \text{tr } A^j$ as before, then we may let

$$10) \quad \tau^\mu = t_1^{m_1} t_2^{m_2} \dots,$$

a product which thus has all factors but a finite number equal to unity.

Each partition μ with $q(\mu) = q$ labels a conjugate class C in the group S_q containing all permutations of q elements, namely that class in which all permutations may be factored into disjoint cycles of which m_1 have length 1, m_2 have length 2, and so forth. The parity of all elements in C_μ is easily shown to be $\text{sgn}(\mu) = (-1)^{p(\mu)}$, and the number of elements in C_μ is well-known to be⁽¹²⁾

$$11) \quad c(\mu) = q! / m_1! m_2! \dots 1^{m_1} 2^{m_2} \dots$$

a quotient whose denominator has all factors but a finite number equal to unity. However, the general expression of the set (3) can then be written⁽¹³⁾

$$12) \quad i! s_i = \sum_{q(\mu)=i} \text{sgn}(\mu) c(\mu) \tau^\mu$$

(12) See 3.6 of Reference 1 or IV.4 of Reference 2.

(13) See 6.2 of Reference 1 or (4.24) of Reference 2.

and the form (1) of the Cayley-Hamilton theorem can be rewritten

$$13) \quad \sum_{i=0}^n (-1)^i A^{n-i} \sum_{q(\mu)=i} \text{sgn}(\mu) c(\mu) \tau^\mu(A) / i! = C.$$

Now we need only completely polarize this equation, noting that it is homogeneous of degree n in A ; that is, we need only replace the n equal variables A by all permutations of n distinct variables A_1, \dots, A_n , and put the sum of all such expressions equal to zero. For each i the corresponding term in (13) then yields $n!$ terms, of which we may collect all those terms such that $A_{\pi(1)}, \dots, A_{\pi(i)}$, in any order, appear in the inner sum, and $A_{\pi(i+1)}, \dots, A_{\pi(n)}$, in any order, appear in the outer sum. Thus for each i we obtain a sum over the $\binom{n}{i}$ ways to select a subset of i elements from $\{A_1, \dots, A_n\}$, with each summand of the form

$$14) \quad (-1)^i [\sum \text{all permutations of } A_{\pi(i+1)} \dots A_{\pi(n)}] \text{coeff.}(A_{\pi(1)}, \dots, A_{\pi(i)})$$

and to find the coefficient for each selection we need only replace the i equal variables A by all permutations of i distinct variables $A_{\pi(1)}, \dots, A_{\pi(i)}$ in

$$15) \quad \sum_{q(\mu)=i} \text{sgn}(\mu) (\text{tr } A)^{m_1} \dots (\text{tr } A^i)^{m_i} / m_1! \dots m_i! 1^{m_1} \dots i^{m_i}$$

and take the sum of all such expressions.

But for each μ in the sum (15) many of the resulting $i!$ terms are equal; in particular, for each j the m_j traces of products containing j factors may be permuted in all $m_j!$ ways without changing the result, and in each of these m_j traces the j factors may be cycled in j ways without changing the result. Since the various terms are otherwise distinct, the order of their

degeneracy is precisely the denominator associated with the given μ in the sum (15), and thus the coefficient is precisely the sum, over all partitions μ of i and all essentially distinct permutations of the $A_{\pi(j)}$, of terms

$$16) \quad \text{sgn}(\mu) \prod_{j=1}^{m_1} \text{tr}(A_{\pi(j)}) \prod_{j=1}^{m_2} \text{tr}(A_{\pi(m_1+2j-1)}, A_{\pi(m_1+2j)}) \dots$$

In summary, the generalized Cayley-Hamilton theorem in n dimensions asserts the vanishing of the sum for $i = 0, \dots, n$ of all essentially distinct terms of the form (14), in which the coefficient is the sum for $q(\mu) = i$ of all essentially distinct terms of the form (16). Furthermore, the coefficients are independent of n , and have the forms given in equation (8) for $i = 1, 2, 3$; finally, by the principle just stated, the coefficient for $i = 4$ has the form

$$17) \quad \begin{aligned} & \text{trAtrBtrCtrD} - \text{trAtrBtrCD} - \text{trAtrCtrBD} - \text{trAtrDtrBC} - \text{trBtrCtrAD} \\ & - \text{trBtrDtrAC} - \text{trCtrDtrAB} + \text{trA}(\text{trBCD} + \text{trDCB}) + \text{trB}(\text{trACD} + \text{trDCA}) \\ & + \text{trC}(\text{trABD} + \text{trDBA}) + \text{trD}(\text{trABC} + \text{trCBA}) + \text{trABtrCD} + \text{trACtrBD} \\ & + \text{trADtrBC} - \text{trABCD} - \text{trABDC} - \text{trACBD} - \text{trACDB} - \text{trADBC} - \text{trADCB} \end{aligned}$$

Since the polarization process d_{BA} can be defined over any field of characteristic zero⁽¹⁴⁾, these results are all valid over any such field. Indeed since the coefficients are simpler in the completely polarized equations, these results may well be provable directly in n variables, rather than through (13). However, this discussion indicates the explicit form of the desired relation, and thus reduces the labor of deriving it to merely that of writing it down.

(14) See p. 4 of Reference 12.

Acknowledgement

I should like to thank Professor R. S. Rivlin for having suggested this question and the explicit form of answer desired, and for having criticized the preliminary drafts of this paper.

The work described in this paper was carried out partly under Contract Nonr 562(40) with the Office of Naval Research and ARPA Contract Sd. 86 with the Advanced Research Projects Agency.

References

1. D. E. Littlewood, The Theory of Group Characters, Oxford 1940
2. F. D. Murnaghan, The Theory of Group Representations, Johns Hopkins 1938
3. A. C. Pipkin and R. S. Rivlin, Arch. Rat'l Mech. Anal. 4, 129 (1959)
4. M. Reiner, Amer. J. Math. 67, 350 (1945)
5. R. S. Rivlin, J. Rat'l Mech. Anal. 4, 681 (1955)
6. G. F. Smith, Arch. Rat'l Mech. Anal. 5, 382 (1960)
7. G. F. Smith and R. S. Rivlin, Arch. Rat'l Math. Anal. 1, 107 (1957)
8. A. J. M. Spencer, Arch. Rat'l Math. Anal. 7, 64 (1961)
9. A. J. M. Spencer and R. S. Rivlin, Arch. Rat'l Math. Anal. 2, 309 (1959)
10. A. J. M. Spencer and R. S. Rivlin, Arch. Rat'l Math. Anal. 2, 435 (1959)
11. A. J. M. Spencer and R. S. Rivlin, Arch. Rat'l Math. Anal. 4, 214 (1960)
12. H. Weyl, The Classical Groups, Princeton 1946

Unclassified

Security Classification

DOCUMENT CONTROL DATA - R&D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) Division of Applied Mathematics Brown University Providence, R. I. 02912		2a. REPORT SECURITY CLASSIFICATION unclassified	
		2b. GROUP ---	
3. REPORT TITLE The Generalized Cayley-Hamilton Theorem in n Dimensions			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) Technical Report			
5. AUTHOR(S) (Last name, first name, initial) Lew, John S.			
6. REPORT DATE March 1966		7a. TOTAL NO. OF PAGES 7	7b. NO. OF REFS 12
8a. CONTRACT OR GRANT NO. Nonr 562(40) <u>and</u> ARPA		9a. ORIGINATOR'S REPORT NUMBER(S) Technical Report No. 6 (ONR) Technical Report No. AM-20 (ARPA)	
b. PROJECT NO. NR-064-4061/3-4-65 <u>and</u> Sd-86		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report) See 9a	
c.			
d.			
10. AVAILABILITY/LIMITATION NOTICES -----			
11. SUPPLEMENTARY NOTES -----		12. SPONSORING MILITARY ACTIVITY Office of Naval Research <u>and</u> Advanced Research Projects Agency	
13. ABSTRACT For any positive integer n , this paper derives the explicit form of the identity in n matrices, each $n \times n$, which is obtained by complete polarization of the usual Cayley/Hamilton theorem, and which is used repeatedly, for $n = 2$ or 3 , in the determination of an integrity basis for symmetric tensors.			

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Cayley-Hamilton theorem n matrix variables integrity basis for tensors						

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through _____."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through _____."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through _____."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.